

New composition and method for the treatment of dysglucaemia

The present invention concerns a method for long term prevention and/or treatment of dysglucaemia and/or for improved glycaemic control, for example the long term prevention of nocturnal and morning hypoglycaemia in diabetic patients, a composition for this purpose and
5 a method for its production.

Cross Reference to Related Applications

This application claims the benefit of U.S. Provisional Application No. 60/243,072, filed October 25, 2000 and Swedish Application No. SE 0003877-8 filed October 25, 2000.

Background of the invention

10 Dysglucaemia is an overall definition, which in the present description and claims is meant to comprise all irregularities in blood glucose level in humans, both healthy persons and persons with a diagnosed disease, irrespective of these irregularities being chronic or temporary, caused by a metabolic disorder, a disease or by physical exercise, bad nutritional habits, medical treatment such as surgery, and desired or undesired pharmaceutical or chemical
15 influences, such as pharmaceutical treatments or substance abuse.

Diabetes mellitus is a complex disorder of the carbohydrate, fat, and protein metabolism that is primarily a result of a relative or complete lack of insulin secretion by the beta cells of the pancreas or of defects of the insulin receptors. The various forms of diabetes are divided in several categories, the two most frequent being juvenile-onset diabetes or Type I insulin-
20 dependent diabetes mellitus (IDDM) and adult-onset diabetes or Type II non-insulin-dependent diabetes mellitus (NIDDM). Both diseases, even when correctly diagnosed and medicated, require life-long medication, good patient compliance, a careful diet and frequent medical observation to avoid potentially serious sequelae.

One problem, frequently encountered by patients suffering from diabetes, is the nocturnal
25 drop in blood glucose levels, hypoglycaemia, in mild cases resulting in morning dizziness and/or nausea. Occasionally the blood glucose level sinks so low during the night, or early in the morning, that the state of hypoglycaemia becomes severe, leading to unconsciousness or convulsions. Importantly, severe hypoglycaemia is more likely to occur at night, when the patient is asleep, rather than during the day, when the patient can feel the onset of

hypoglycaemia and prevent it by eating carbohydrates, e.g. a lump of sugar or specific glucose tablets, energy gels or bars, marketed for diabetic patients.

Moreover, both mild (a blood glucose level about 2.4 – 4.0 mmol/l) and severe (less than 2.4 mmol/l) hypoglycaemia predisposes the patient to a condition known as hypoglycaemic
5 unawareness, which in turn means that hypoglycaemia can occur more frequently and at any time of the day, due to attenuation of the typical warning symptoms of a declining blood glucose level, e.g. hunger, perspiration etc.

Notably, the incidence of hypoglycaemia is rapidly increasing as diabetic patients seek better control of their blood glucose levels, in order to avoid hyperglycaemia. An active life,
10 involving exercise as prescribed for diabetics, also adds to the risk of hypoglycaemia unless the patient carefully controls his/her carbohydrate intake, insulin dosage and regularly monitors their blood glucose level.

There is presently no satisfactory therapeutic regime for the prevention of nocturnal and morning hypoglycaemia. Patients suffering from diabetes are often recommended to eat a
15 light meal before bedtime, e.g. potato chips or cereals. This results in an almost instant peak in blood glucose, followed by a more or less rapid decline during the night. In order to guarantee a sufficient blood glucose level throughout the night, the initial level must be very high, bordering to the unhealthy. When attempting this kind of crude self-medication, the patients have difficulty finding the optimal dose and mix of carbohydrates, and over-eating
20 tends to be the result. There is currently no medication available with an appropriate kinetic profile to counteract the effect of the night-time intermediate-acting insulin.

Prior art

A method of treating diabetic hypoglycaemia by administration of uncooked cornstarch is described in WO95/24906. According to this application, published in September 1995, the
25 patient is given about 0.1 to 1.0 g starch per kg body weight, preferably about 0.25 to 0.5 g per kg body weight. Different dosage formulations are suggested, including a suspension of starch in milk, sustained release tablets and a snack bar, containing a total of 30 g carbohydrates, but having about 1/2 to 1/4 of the carbohydrate in the form of uncooked cornstarch. Although containing "slow" carbohydrates in the form of cornstarch, this product
30 contains considerable amounts of free sugar and fats, resulting both in an initially very high

blood glucose peak, and in the intake of unnecessary calories in the form of fats and surplus carbohydrates. Further information concerning the sustained release tablets is not given, apart from a reference to well known techniques of tablet formulation.

U.S. 5,605,893 (Kaufman, F.) discloses a specific method of treating a diabetic patient and preventing hypoglycaemic episodes, said method consisting of administration to the patient of a therapeutic food composition comprising per serving or unit about 20 – 50 grams of nutrients, including

- about 5 – 15 g of slowly absorbed complex carbohydrate, preferably cornstarch;
- about 7 – 19 g of rapidly absorbed complex carbohydrate;
- 10 - about 5 – 20 g of protein; and
- about 3 – 7 g of fat,

said composition being substantially free from simple sugars.

U.S. 5,843,921 (Kaufman, F.) discloses a therapeutic food composition for treatment of diabetic patients and preventing hypoglycaemic episodes, comprising per serving or unit about 20 – 50 grams of nutrients, including

- about 5 – 15 g of slowly absorbed complex carbohydrate, preferably cornstarch;
- about 7 – 19 g of rapidly absorbed complex carbohydrate;
- about 5 – 20 g of protein; and
- about 3 – 7 g of fat;

20 wherein the amount of simple sugars other than fructose in said composition is less than about 3 g per unit.

U.S. 5,866,555 (Bell, S. J. *et al.*) discloses a diabetic supplement bar for the treatment or prevention of night time hypoglycaemia in a diabetic patient, made by blending simple carbohydrates, proteins, lipids, complex carbohydrates, and any additional additives, and homogenising the mixture into a food bar. Sucrose is presented as the preferred source of simple carbohydrates, whereas uncooked cornstarch is the preferred source of complex carbohydrates. The final fat concentration of the product is high, from 2 to 40 % by weight. The remaining ingredients are:

- about 10 – 60 % by weight sucrose;
- 30 - about 1 – 25 % by weight protein;
- about 1 – 60 % by weight complex carbohydrate.

The high fat content of the above products accounts – together with the naturally slow degradation of uncooked cornstarch – for the delayed glucose release, as fat delays gastric emptying, thereby slowing the rate at which nutrients enter the intestines and become digested.

- 5 Another product, available on the market under the trade mark NiteBite® (Optim Nutrition Inc.) contains three sources of glucose: sucrose, protein and uncooked cornstarch. These components are digested more or less consecutively and are claimed to deliver glucose into the blood during a period of 6 hours or more.

10 U.S. 5,545,410 discloses the coating of a core of a carbohydrate in a food or pharmaceutical grade coating such as stearic acid, hydrogenated or partially hydrogenated oils, calcium stearate, stearyl alcohol, wax etc. The claims of U.S. 5,545,410 mention sucrose, glucose, lactose, dextrans, monosaccharides, disaccharides, oligosaccharides, and polysaccharides, as well as pregelatinized starches, raw starches, and modified starches. A limitation is however
15 individually coated particles or granulates thereof, the properties of the coating regulating the release of the core compound.

The related patent U.S. 5,360,614 discloses the above coated carbohydrate cores, further defining their size to the interval of 75 – 500 μm , and the coating as providing for the release of substantially all said metabolizable carbohydrate into the digestive system for one-half to
20 four hours after ingestion.

The prior art compositions fail to provide an entirely satisfactory solution for long term treatments when all effects and consequences are taken into account. The sucrose included in some preparations can lead to initial hyperglycaemia and may additionally contribute to caries and tooth decay. The proteins, and in particular the fat included in some of the above
25 preparations provide unnecessary additional calories to patients, who in many cases already battle with weight problems.

The objective of the present invention is thus to accurately and reliably regulate blood glucose levels in humans and prevent dysglucaemia, in particular long term prevention of dysglucaemia. This objective can be divided in the prevention of dysglucaemia in diabetic
30 patients, and in particular the prevention of nocturnal dysglucaemia in diabetic patients.

Another related objective is to prevent nocturnal hypoglycaemia in diabetic patients and in particular in type 1 and type 2 diabetics on insulin medication.

A further problem is how to prevent and/or treat dysglucaemia in both healthy subjects, such as athletes, and in unhealthy, such as patients undergoing surgery, chemotherapy etc, and in particular diabetic patients undergoing medical treatment.

Further, it is desirable to make available a simple and yet reliable way of improving the glycaemic control or stabilising the oscillating blood sugar levels typical for diabetic patients, and avoiding both the high and low blood sugar levels, both conditions injurious to the health, especially when occurring repeatedly or for prolonged periods of time.

10 It is particularly desirable to make available a method and composition giving a controlled, delayed enzymatic digestion of the starch and preferably a substantially linear glucose release curve and thus guaranteeing a pre-determined, stable and sufficient blood sugar level during at least 5 hours, preferably about 8 hours.

A practical problem encountered in pursuing the above objectives is how to make available a cornstarch composition having an agreeable taste and texture, suitable for daily consumption and life long treatment. This is not without significance, as a tasty product advances good compliance.

Summary of the invention

The present inventors have surprisingly found that the conversion of starch into sugar and in particular the enzymatic conversion of cornstarch into glucose, can be regulated or delayed in a controlled manner and adapted to the metabolism of the patient, without relying on the delaying effect of fat. Thus the blood sugar level can be held at a desired level, e.g. a level accurately adjusted to the metabolic needs of the patient, avoiding both peaks with the associated risk of hyperglycaemia, and low levels, and the corresponding risk of hypoglycaemia. Importantly, this can now be achieved without the administration of unnecessary surplus calories. Further, the present invention makes available a tasty and palatable product consisting substantially of native cornstarch.

The problems associated with the prior art products are solved, and the above objectives of the invention are achieved by a novel composition and a method comprising administering to

the patient in question a starch product with minimised available surface area, such as a granulated and/or partially encapsulated starch, preferably native cornstarch. The method of treatment, the pharmaceutical composition itself and the process for its production are as disclosed in the attached claims.

5

Short description of the drawings

The present invention will be described in closer detail in the following description, examples and enclosed drawings, in which

Fig. 1 shows the *in vitro* degradation profiles for different compositions according to the invention, compared to the profile for untreated cornstarch;

10 Fig. 2 shows the average fasting and postprandial blood glucose levels in healthy, fasting volunteers, after ingestion of 4 different compositions according to the invention;

Fig. 3 shows the correspondence between rates of digestion / degradation as measured by *in vivo* and *in vitro* techniques;

15 Fig. 4 shows the *in vivo* blood glucose contribution in a diabetic patient after intake of a light meal compared to native cornstarch; and

Fig. 5 shows the *in vivo* blood glucose curve of a diabetic patient having taken the inventive composition, compared to control, excipients only, during ongoing insulin infusion according to the glucose clamp technique. The upper curve thus shows the contribution of the inventive composition.

20

Description

Starch is the main storage polysaccharide in plants, an important source of carbohydrates and an ingredient in food. More than half of the carbohydrates ingested by humans is starch.

Starch also has many technical uses, accounting for large volumes of the starch produced.

25 Examples include the use as a sizing agent in paper industry, a raw material and/or additive in the production of plastics, textiles etc. and as a carrier and bulk agent in pharmaceutical industry. Starch is also used as a non-adherence agent for surgical gloves. Apart from these purely technical uses, it is not known to the present inventors that native starch as such would

have a purely therapeutic, internal use, nor that it would have constituted the active ingredient in a pharmaceutical composition, such as a tablet as disclosed in the present description and claims.

5 Pure starch, although theoretically a good source of glucose and free from surplus calories, is practically impossible to ingest. The starch powder itself lacks taste and feels extremely dry, chalky or sandy in the mouth. A suspension on the other hand, e.g. a suspension in water, tends to sediment quickly, and has a disagreeable taste and texture.

Cornstarch consists of granules sized 2 – 32 μm , mainly comprising two components, amylose and amylopectin. Amylose has a linear structure while amylopectin is branched.
10 Both amylose and amylopectin consist of α -(1,4)-linked glucose residues while amylopectin also has α -(1,6)-linked glucose residues. The starch granules are insoluble in cold water and swell in warm. The swelling is reversible until the temperature reaches about 55 to 65 $^{\circ}\text{C}$. At this temperature the starch granules gelatinise and lose their crystalline structure.

The degradation of starch is catalysed by α -amylase, which in humans is present in the saliva
15 and in the small intestine. The digestibility of starch, both *in vivo* and *in vitro* depends on the source of starch as well as its pre-treatment (e.g. native, fine / coarse, gelatinised or chemically modified). In the present description, claims and examples, the term “native starch” is used to define starch that has not been subjected to heat-treatment or chemical treatment. The term “native starch” thus comprises both the vegetable and/or plant seeds,
20 kernels or grains, as well as mechanically treated fractions, such as the milled and sieved product, granules and flour.

The present inventors have now surprisingly found, that the enzymatic degradation of the starch *in vivo* can be regulated in an accurate and repeatable manner by minimising the surface area available to enzymatic action, preferably by granulating the starch granules with
25 a substance, resulting in aggregated granules being at least partially encapsulated in the substance.

Suitable substances are non-toxic substances, suitable for ingestion, such as substances generally recognised as safe (GRAS) and approved for use in pharmaceutical applications and/or in food products. A non-exclusive list of suitable substances includes polymers such as
30 gum arabicum, potassium alginate, guar gum, methyl cellulose, ethyl cellulose; liquid oils,

liquid and hard fats and waxes, such as paraffin, hydrogenated cottonseed oil, beeswax, and carnauba wax.

Tests conducted *in vitro*, have shown that a controlled enzymatic digestion and a substantially linear release profile is achieved when the amount of reducing sugars is plotted against time.

- 5 Test conducted *in vivo*, using healthy volunteers, have shown that a modulated release profile is achieved. This makes it possible to achieve a long term delivery of reducing sugars, adjusted in level and duration to the metabolic needs of the patient. An additional *in vivo* test, conducted on a diabetic patient, confirms the performance of the inventive composition and the feasibility of the inventive method. This test was performed after the priority date, but
10 before the international filing date.

- The present invention makes available a method and composition for the long term prevention and/or treatment of dysglucaemia, e.g. the prevention of nocturnal and/or morning hypoglycaemia in patients suffering from diabetes, including both IDDM and NIDDM, wherein a predetermined amount of starch is administered to the patient in granulated and at
15 least partially encapsulated form, most preferably in the form of tablets, which granulation delays the enzymatic degradation of the starch into reducing sugars, such as glucose. The starch is preferably native cornstarch.

- The present invention also makes available a method and composition for the prevention of dysglucaemia in situations, where the blood glucose level is disturbed or altered by exercise, pharmaceutical or surgical therapy, by a disease or a syndrome, involving multiple diseases or
20 metabolic disorders. Examples include athletes, patients weakened by chemotherapy, fasting patients and patients suffering from diseases or disorders disturbing or altering the sugar metabolism, or patients undergoing treatment of such and other diseases or disorders. The method is characterised in that a predetermined amount of starch is administered to the patient
25 in granulated and at least partially encapsulated form, preferably in the form of tablets, which granulation delays the enzymatic degradation of the starch into reducing sugars, such as glucose. The starch is preferably native cornstarch.

- The present invention also makes available a method and composition for improved glycaemic control, i.e. for stabilising the blood sugar levels and avoiding the oscillation
30 between unhealthy high and low blood sugar levels. Lack of glycaemic control is associated

- optionally about 0.1 - 15 % by weight heat treated cornstarch in the form of flakes;
- about 5 – 25 % by weight low calorie sweetener, e.g. isomalt, sorbitol, xylitol, or the like, preferably isomalt.

According to another embodiment, the inventive composition for controlled release of glucose
5 comprises the following ingredients:

- about 60 - 90 % by weight native cornstarch encapsulated in a substance,
- optionally about 0.1 - 15 % by weight heat treated cornstarch in the form of flakes;
- about 0.1 - 10 %, preferably 0.1 – 5 % by weight unsaturated fat;
- 0.01 – 25 % by weight low calorie sweetener, e.g. isomalt, sorbitol, xylitol, saccharin, or
10 the like, preferably isomalt.

Preferably the heat treated flakes have a size in the interval of 0.5 to 1.0 mm and the unsaturated fat is olive oil. The composition may further contain optional additives, such as additives giving the final product its colour and taste.

The heat treated starch may be added in the form of baked flakes, which give the product a
15 crispy texture and a pleasant taste. Flakes suitable for this purpose are made of a mixture of cornstarch, water, sodium chloride and sweetener. The flakes are heated above gelatinisation temperature and subsequently dried at about 90 °C. The dried flakes are milled and sieved. Flakes in the interval of about 0.5 – 1.0 mm are then mixed with the granulated and at least partially encapsulated cornstarch.

20 Preferably the substance which encapsulates the starch is chosen among non-toxic substances, suitable for ingestion, such as substances generally recognised as safe (GRAS) and approved for use in pharmaceutical applications and/or in food products. A non-exclusive list of suitable substances includes polymers such as gum arabicum, potassium alginate, guar gum, methyl cellulose, ethyl cellulose; liquid oils, liquid and hard fats and waxes, such as paraffin,
25 hydrogenated cottonseed oil, beeswax, and carnauba wax.

Preferably the substance is chosen among guar gum and ethyl cellulose, most preferably ethyl cellulose.

inter alia with microvascular damage, such as diabetic retinopathy, and diabetic ketoacidosis or so called diabetic coma.

The present invention further makes available a composition for controlled release of reducing sugars, e.g. glucose, wherein said composition contains granulated and at least partially encapsulated starch, a low calorie sweetener and optionally also unsaturated fat. The starch is preferably native cornstarch.

According to one embodiment of the invention, the native starch is granulated with and at least partially encapsulated in a substance chosen among non-toxic substances, suitable for ingestion, such as substances generally recognised as safe (GRAS) and approved for use in pharmaceutical applications and/or in food products. A non-exclusive list of suitable substances includes polymers such as gum arabicum, potassium alginate, guar gum, methyl cellulose, ethyl cellulose; liquid oils, liquid and hard fats and waxes, such as paraffin, hydrogenated cottonseed oil, beeswax, and carnauba wax.

Preferably the substance is chosen among guar gum and ethyl cellulose, most preferably ethyl cellulose.

According to the invention, the enzymatic degradation is delayed to an extent resulting in a controlled and substantially linear release of reducing sugars, e.g. glucose, accurately adapted to the metabolism of the patient for more than 4 hours, preferably more than 6 hours, most preferably about 8 hours or longer.

According to one embodiment, the composition comprises starch in two forms; a first amount of native starch, and a second amount of heat treated starch. Preferably the second amount of heat treated starch is about 0.1 to 15 % by weight of the total amount of starch, preferably about 5 % by weight.

According to a preferred embodiment the heat treated and un-encapsulated starch is in the form of flakes of baked starch having a size in the interval of about 0.5 – 1.0 mm.

According to one embodiment, the inventive composition for controlled release of glucose comprises the following ingredients:

- about 60 - 90 % by weight native cornstarch encapsulated in a substance,

According to a preferred embodiment, an organic acid is added to the inventive composition. The organic acid is preferably one of ascorbic acid, tartaric or citric acid.

The inventive composition may further comprise minor amounts of technical additives, such as conventional tableting agents, lubricants and glidants.

5 The present invention discloses a method for production of a composition for the controlled delayed degradation of starch, wherein the method comprises the following steps

- granulation and at least partial encapsulation of native cornstarch in a substance which delays the degradation,
- wet sieving and drying of the encapsulated granules,
- 10 - adding a low calorie sweetener and an organic acid, and
- pressing the mixture into tablets.

As one embodiment, the present invention further discloses a method for production of a composition for the delayed degradation of starch, wherein the method comprises the following steps

- 15 - granulation and at least partial encapsulation of native cornstarch in a substance which delays the degradation,
- wet sieving and drying of the encapsulated granules,
- mixing the granules with heat treated flakes of cornstarch,
- adding a low calorie sweetener, an organic acid and unsaturated fat to the mixture, and
- 20 - pressing the mixture into tablets.

The tablets are then packaged according to conventional methods, in a package suitable for storage, delivery and sale. Optionally, the granulate is not pressed into tablets but weighed and packaged according to conventional methods, in a package suitable for storage, delivery and sale.

The substance which delays the degradation is chosen among non-toxic substances, suitable for ingestion, such as substances generally recognised as safe (GRAS) and approved for use in pharmaceutical applications and/or in food products. A non-exclusive list of suitable substances includes polymers such as gum arabicum, potassium alginate, guar gum, methyl cellulose, ethyl cellulose; liquid oils, liquid and hard fats and waxes, such as paraffin, hydrogenated cottonseed oil, beeswax, and carnauba wax. Preferably the substance is chosen among guar gum and ethyl cellulose, most preferably ethyl cellulose.

The heat treated starch may be added in the form of baked flakes, which give the product a crispy texture and a pleasant taste. Flakes suitable for this purpose are made of a mixture of cornstarch, water, sodium chloride and sweetener. The flakes are heated above gelatinisation temperature and subsequently dried at about 90 °C. The dried flakes are milled and sieved. Flakes in the interval of about 0.5 – 1.0 mm are then mixed with the granulated and at least partially encapsulated cornstarch before tablet pressing.

According to a preferred embodiment, an organic acid is added to the mixture before tablet pressing, the organic acid being chosen among ascorbic acid, tartaric acid and citric acid, preferably tartaric acid. An organic acid has the additional advantage of stimulating the secretion of saliva. This is especially advantageous as pure starch and products with a high starch content have a dry and chalky or sandy mouth feel and bland or even disagreeable taste.

The product according to the present invention is delivered in the form of a granulate, or preferably in the form of pressed cakes or tablets. The pressed cakes or tablets are preferably produced with grooves or notches for easy division into even sized fractions.

According to one embodiment of the invention, a saturated or preferably unsaturated fat is added to the granulate before tablet pressing. Preferably an unsaturated fat is used, and most preferably olive oil, added in about 0.1 – 5 % by weight for tabletted products, and 0.1 – 10 % by weight for granulates.

According to one embodiment of the invention a low calorie sweetener is added, and preferably saccharin, isomalt or xylitol is used as the sweetener. Xylitol imparts a cooling sensation because of its endothermic dissolution.

A specific advantage of the method and composition of the present invention is that practically all of the starch, contained in the ingested dose, is converted to reducing sugars, mainly glucose. Thus the amount of glucose can be accurately calculated and the dose optimised for each patient, e.g. by producing tablets of different size or by ordinating the patient to take a prescribed number of tablets. Further, a minimum of surplus calories are administered to the patient.

A particular advantage of the present composition and method is that the release rate and the content of reducing sugars can be accurately controlled and adjusted to the needs of the specific patient group, specific application or medical situation, and adjusted to the metabolism of the patient or patient. A method and product according to the invention makes possible an exact and reliable dosage and ease of use.

Another advantage of the composition according to the present invention is that undigested starch is prevented from reaching the colon, where it would be digested by bacteria, resulting in the formation of gas, especially in the colon.

Another advantage is that – for the first time known to the inventors – a substantially pure starch composition has been formulated as a product with agreeable taste and mouthfeel. As touched upon above, the sensory qualities of a medical product have importance for compliance.

Examples

In vitro degradation tests

The tested formulations were prepared using a high shear mixer (Donsmark QMM-II) and tablets pressed with a hydraulic single press (Compac DP6-B2) or with a rotary tablet press (Korsch Pharmapress PH-106). All formulations were based on native cornstarch (Maizena, Bestfoods Nordic AB). Cornstarch and different excipients were dry mixed in a granulator and agglomerated with water or ethanol as granulation fluid, depending on the solubility of the granulation substance used. The dry granulate was pressed into tablets.

In order to study the enzymatic degradation of starch *in vitro*, a novel analytical method was developed, and is the object of co-pending application PCT/SE01/02298. According to this method, the degradation resistance of starch by the action of α -amylase is measured as the

concentration free sugar in a starch suspension incubated at 37 °C after the addition of enzyme. Samples are taken at regular intervals and the reducing sugars, e.g. glucose and maltose, are reacted with a reagent consisting of a filtered 3,5-dinitro salicylate solution in aqueous NaOH. The formed colour is determined spectrophotometrically by scanning over
5 the wave length interval of 450 – 500 nm, detecting the absorption maximum. Enzymatic degradation is then plotted as sugar concentration as a function of the incubation time.

Example 1: Cornstarch granulated with potassium alginate (10 %) compared to free cornstarch

Method of production: Native cornstarch (Maizena, Bestfoods Nordic AB) was mixed in a
10 high shear mixer with potassium alginate (10 % by weight, Food grade, Danisco) and water added as granulation fluid. The granules were wet sieved through a 1 mm sieve and dried in a forced hot air oven (Fermaks) at 35 – 40 °C. The dried granulate was sieved through a 1 mm sieve and collected on a 0.5 mm sieve. A small amount of fat was added [5, 8 and 10 % olive oil (food grade, Zeta), or 5, 10 and 20 % Akosupp 10 (Karlshamns AB)]. The granulate and/or
15 tablets were tested *in vitro* according to the method described below, and *in vivo* according to the method described further below, in connection with the *in vivo* tests.

Method of analysis: A reagent was made by dissolving 3,5-dinitro salicylate (2.00 g, Aldrich) in aqueous NaOH (70 ml, 1 M). Optionally, the mixture is heated in order to expedite the formation of a clear solution. Upon cooling, water is added to 100 ml. The reagent solution is
20 stored in a dark place and filtered through a 0.45 µm filter before use, in order to remove possible precipitates.

The reagent solution was added in equal amounts (2 ml) in test tubes marked "control", "zero", "5 min", "10 min", "20 min", "30 min", "45 min", "1 h", "1.5 h", "2 h", "2.5 h", "3 h", "3.5 h", and "4 h". The test tubes were placed in an ice bath awaiting the analysis.

25 A buffer solution (pH 6.6) was made by mixing KH_2PO_4 (250.0 ml, 0.20 M, Sigma) and NaOH (89.0 ml, 0.20 M) and adding water to a total volume of 1000 ml. NaCl (0.58 g, Riedel-de Haën) was then added to produce a chloride concentration of 0.01 M.

A defined amount of starch to be investigated is suspended in the above buffer and placed in the degradation bath. The degradation bath is kept at a temperature of 37 °C ± 0.5 °C and
30 stirred at a speed of 50 rpm.

- An amount corresponding to 15 000 IU α -amylase (Type VI-B from porcine pancreas, Sigma) is measured and suspended in buffer. Before addition of the enzyme solution, a sample of the degradation bath is taken in order to determine the sugar concentration at "time zero". The sample is filtered through a 0.8 μ m filter and an aliquot (2 ml) is pipetted to the
- 5 test tube marked "zero". The same filter can be used throughout the series. The sample is boiled momentarily (5 min) and placed in an ice-bath. Following this, the enzyme solution is added to the degradation bath and the time registered. Samples are then taken at predetermined intervals, such as the times indicated on the test tubes. The control is prepared by boiling reagent (2 ml) and water (2 ml) during 5 min and placing the sample in an ice-bath.
- 10 For each sample, the absorption is scanned in the interval 450 – 500 nm and the peak height registered for each absorption maximum. In order to determine the concentration free sugars (FS0) in the native starch, the absorbance of the sample "zero" is measured against a background of water and reagent, the control sample. Both samples and control are diluted by adding 11.6 ml water to 400 μ l sample. The reacted and diluted sample solution is not stable
- 15 (the reading falling 0.1 to 0.2 absorbancy units during 3 hours) so all samples are diluted slightly prior to the UV-spectrophotometric analysis.

Example 2: Cornstarch granulated with potassium alginate (20 %) compared to free cornstarch

- Native cornstarch was granulated with 20 % by weight potassium alginate and analysed as
- 20 described in Example 1.

Example 3: Cornstarch granulated with guar gum (20 %) compared to free cornstarch

Native cornstarch was granulated with 20 % by weight guar gum (Scanpharm A/S) and analysed as described in Example 1.

Example 4: Cornstarch granulated with ethyl cellulose compared to free cornstarch

- 25 Native cornstarch was granulated with 18 % by weight ethyl cellulose (Dow Chemical Co.) and analysed as described in Example 1.

The results of the *in vitro* degradation tests are shown in Fig. 1, where the values for cornstarch encapsulated in ethyl cellulose (10 %) are marked with the symbol (O), cornstarch

encapsulated in ethyl cellulose (20 %) with (Δ), cornstarch encapsulated in guar gum (20 %) being marked with (9) and the values for untreated cornstarch being marked (6).

In vivo tests

- 5 The blood glucose response was measured by the standard technique in 4 healthy, lean volunteers (age 35 to 45 years) with normal glucose tolerance. According to the “golden standard” of this technique, each substance was studied twice in each volunteer, and the mean value was calculated. The substances were tested in randomised order, at least one day apart. Moreover, the testing was performed under strictly standardised conditions. The subjects
- 10 came to the laboratory in the morning, fasted for 10 hours. Physical activity was avoided right before and during the test. The test subjects were allowed to drink about 2 dl liquid, free from carbohydrates (water, tea or coffee) twice during the test; at 0 and 3 hours.

- The capillary blood glucose level was determined in capillary blood samples (obtained by finger pricking) using a Glucometer DEX (Bayer Diagnostica AB) following the standard
- 15 procedures for glucose measurements. At baseline, three consecutive blood glucose determinations were performed to ensure a stable baseline at time 0 hours. Thereafter the test substance (20.0 g) was ingested together with a standardised amount of water within 5 minutes. All liquids were carefully weighed and the same amounts ingested at each occasion to avoid variations in transit time through the gastrointestinal tract. The blood glucose
- 20 determination was repeated at 0.5, 1.0, 1.5, 2.0, 3.0, and 4.0 hours. The average results for the test subjects are shown in Fig. 2. The correspondence between rates of digestion as measured by *in vivo* (4 h blood glucose response, ((mmol/l min)*100) and *in vitro* (as described in Example 1) techniques is shown in Fig. 3.

Example 5: In vivo effect of cornstarch granulated with ethyl cellulose (10 %)

- 25 The test subjects measured their initial blood glucose after 10 hours of fasting, whereupon they ingested an exactly measured amount of a formulation of cornstarch granulated with and partially encapsulated in ethyl cellulose (10 %), manufactured according to the method presented in Example 1 above. Blood glucose was then measured at 0.5, 1.0, 1.5, 2.0, 3.0, and
- 30 4.0 hours. The average result for all test subjects are presented in Fig. 2, the values for cornstarch encapsulated in ethyl cellulose (10 %) being marked with (O).

Example 6: In vivo effect of cornstarch granulated with ethyl cellulose (20 %)

The test subjects measured their initial blood glucose after 10 hours of fasting, whereupon they ingested an exactly measured amount of a formulation of cornstarch encapsulated in ethyl cellulose (20 %), manufactured according to the method presented in Example 1 above.

- 5 Blood glucose was then measured at 0.5, 1.0, 1.5, 2.0, 3.0, and 4.0 hours. The average result for all test subjects are presented in Fig. 2, the values for cornstarch encapsulated in ethyl cellulose (20 %) being marked with (Δ).

Example 7: In vivo effect of cornstarch granulated with guar gum (20 %)

- 10 The test subjects measured their initial blood glucose after 10 hours of fasting, whereupon they ingested a formulation of cornstarch encapsulated in guar gum (20 %) manufactured according to the method presented in Example 1 above. Blood glucose was then measured at 0.5, 1.0, 1.5, 2.0, 3.0 and 4.0 hours. The average result for all test subjects are presented in Fig. 2, the values for cornstarch encapsulated in guar gum (20 %) being marked with (9).

15 **Example 8: In vivo effect of untreated cornstarch**

The test subjects measured their initial blood glucose after 10 hours of fasting, whereupon they ingested native, untreated cornstarch (Maizena, Bestfoods Nordic AB). Blood glucose was then measured at 0.5, 1.0, 1.5, 2.0, 3.0, and 4.0 hours. The average result for all test subjects are presented in Fig. 2, the values for untreated cornstarch being marked (6).

- 20 The *in vivo* results clearly show that a marked effect is achieved with the formulations according to the invention, compared to untreated cornstarch. Most notably, the initial glucose peak appearing at 0.5, 1.0 and 1.5 hours after ingestion of untreated cornstarch, is entirely absent for the guar gum encapsulated cornstarch.

Clinical trials

25 **Example 10: Comparative tests performed with diabetic patients**

Study design: The 16 patients in the study will receive the inventive composition in tablet form (MM005) in different single doses, 5, 10 and 20 g, respectively, at three different occasions and the control (excipients only) at one occasion.

Establishment of baseline (control occasion): The subjects come to the laboratory in the morning in the fasting state and without having taken their regular morning insulin dose. With the help of a slow i.v. infusion of insulin, combined with a glucose infusion, the blood glucose level will be stabilised at 5.5 to 6.5 mmol/L. The insulin will be administered by an infusion rate, aiming at giving a blood insulin concentration of 15 – 20 mU/L. The glucose concentration will be locked by customary clamp technique, where blood sugar is measured every 5th minute for 1 hour and the glucose infusion rate is adjusted if necessary to give the desired blood glucose concentration. Thereafter the control medication is given and the glucose clamp continued for 6 hours. The amount of glucose administered during each hour will be used at the three subsequent test occasions. The objective is to establish a reference baseline for evaluation of doses.

Dose testing: The sequence of the subsequent three test occasions, i.e. for the three different doses of the investigational drug, will be randomised. At least one week should pass between test occasions. Every test day will be started by glucose and insulin infusion for 1 hour as described above. Thereafter the investigational drug is administered orally as a single dose and the glucose infusion is continued so that the amount of given glucose during each hour corresponds to the amount given on day 1 (control occasion). During the three test days blood samples are withdrawn every 10th minute during the first 6 hours of the experiment for glucose determination, and also every 60th minute for insulin determination. The subject will be asked to return within 1-4 weeks for a new test.

Dose: The doses are 5 g, 10 g, and 20 g cornstarch, i.e. 3, 6 and 12 tablets. The dose is taken once a day, in the morning, as a single dose.

Drug administration: The formulated cornstarch is administered orally as chewing tablets formulated according to the invention, together with 250 ml of water.

Discussion of Study Design: The aim of this study is to evaluate the increase in glucose levels following intake of three cornstarch doses (5, 10, 20 g) as well as the kinetic profile compared to a control period with inactive medication (excipients only).

This is important since the medication is to be used to prevent hypoglycemia, in particular nocturnal hypoglycemia. This usually occurs at between 02 and 04 in the morning.

There is currently no medication available with an appropriate kinetic profile to counteract an excessive effect of the night-time intermediate-acting insulin. Thus, diabetic subjects usually ingest food when they have measured low glucose levels at bedtime. However, very high early night-time glucose levels are necessary to provide a sufficient "buffer" at 02-04 o'clock in the morning. Thus, a medication which provides such an effect should be an important therapy in diabetes.

The doses chosen are within the range calculated to be required to counteract the effect of an excessive nocturnal insulin dose.

The use of a control medication (excipients only) provides a background which makes it possible to evaluate the glucose-releasing property of the active medication. The number of subjects included in the study has been calculated to give enough power to safely evaluate the glucose-releasing profile.

The subjects should not have any clinically significant manifestations of the specific diabetic microvascular disorders.

The subjects will be given glucose if the blood glucose levels drop below 3.5 mM, and the test will be discontinued on that day.

The first data available from the clinical studies show that the inventive composition is capable of stabilising the blood glucose over a period of at least 6 hours, and that the enzymatic degradation of starch / conversion into glucose exhibits a substantially linear behaviour.

In Fig. 4, the glucose contribution of a light meal (v), is compared to that of native cornstarch (♦). The benefits of native cornstarch are clearly visible from the diagram. The native cornstarch formulation totally lacks the initial glucose peak, and gives a steady contribution still after 8 hours, with the trend of the curve indicating a considerably longer effect than for that accountable to a light meal.

See also Fig. 5 which shows the glucose curve of a diabetic patient having eaten the inventive composition (Δ), compared to control, excipients only (O). The results display a pronounced effect of the inventive composition still after 6 hours, the trend of the curve indicating that the

effect most probably lasts even longer. These tests were performed after the priority date, but before the international filing date.

In organoleptic tests, the test subjects described the formulations according to the present invention as “tasty”, “crispy” and “easy to swallow”.

- 5 Although the invention has been described with regard to its preferred embodiments, which constitute the best mode presently known to the inventors, it should be understood that various changes and modifications as would be obvious to one having the ordinary skill in this art may be made without departing from the scope of the invention as set forth in the claims appended hereto.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186